

Amsterdam University of Applied Sciences

Factors Associated with Step Numbers in Acutely Hospitalized Older Adults: The Hospital-Activities of Daily Living Study

Kolk, Daisy; Aarden, Jesse J. ; MacNeil-Vroomen, Janet L. ; Reichardt, Lucienne A. ; van Seben, Rosanne; van der Schaaf, Marike; van der Esch, Martin; Twisk, Jos W.R. ; Bosch, Jos A. ; Buurman, Bianca M. ; Engelbert, Raoul H.H. ; Hospital-ADL study group

DOI

[10.1016/j.jamda.2020.06.027](https://doi.org/10.1016/j.jamda.2020.06.027)

Publication date

2021

Document Version

Final published version

Published in

Journal of the American Medical Directors Association

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Kolk, D., Aarden, J. J., MacNeil-Vroomen, J. L., Reichardt, L. A., van Seben, R., van der Schaaf, M., van der Esch, M., Twisk, J. W. R., Bosch, J. A., Buurman, B. M., Engelbert, R. H. H., & Hospital-ADL study group (2021). Factors Associated with Step Numbers in Acutely Hospitalized Older Adults: The Hospital-Activities of Daily Living Study. *Journal of the American Medical Directors Association*, 22(2), 425-432.
<https://doi.org/10.1016/j.jamda.2020.06.027>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the library: <https://www.amsterdamuas.com/library/contact/questions>, or send a letter to: University Library (Library of the University of Amsterdam and Amsterdam University of Applied Sciences), Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.



JAMDA

journal homepage: www.jamda.com

Original Study

Factors Associated with Step Numbers in Acutely Hospitalized Older Adults: The Hospital-Activities of Daily Living Study



Daisy Kolk MSc^{a,*,†}, Jesse J. Aarden MSc^{b,c,†}, Janet L. MacNeil-Vroomen PhD^a,
 Lucienne A. Reichardt PhD^{a,d}, Rosanne van Seben PhD^{a,e},
 Marike van der Schaaf PhD^{b,c}, Martin van der Esch PhD^{c,f}, Jos W.R. Twisk PhD^g,
 Jos A. Bosch PhD^{h,i}, Bianca M. Buurman PhD^{a,c},
 Raoul H.H. Engelbert PhD^{b,c}, on behalf of the Hospital-ADL Study Group[‡]

^a Amsterdam UMC, University of Amsterdam, Internal Medicine, Section of Geriatric Medicine, Amsterdam Public Health Research Institute, Amsterdam, Netherlands

^b Amsterdam UMC, University of Amsterdam, Rehabilitation, Amsterdam Movement Sciences, Amsterdam, Netherlands

^c ACHIEVE, Center of Applied Research, Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, Netherlands

^d Statistics Netherlands, Department Health and Care, The Hague, the Netherlands

^e Ecorys Nederland, Social Affairs and Health, Rotterdam, the Netherlands

^f Reade, Center for Rehabilitation and Rheumatology/Amsterdam Rehabilitation Research Center, Amsterdam, the Netherlands

^g Amsterdam UMC, Vrije Universiteit Amsterdam, Epidemiology and Biostatistics, Amsterdam, the Netherlands

^h University of Amsterdam, Department of Clinical Psychology, Amsterdam, the Netherlands

ⁱ Amsterdam UMC, University of Amsterdam, Medical Psychology, Amsterdam, the Netherlands

A B S T R A C T

Keywords:

Accelerometer
 physical performance
 hospitalization
 post-acute care
 postdischarge
 older adults

Objectives: To determine the number of steps taken by older patients in hospital and 1 week after discharge; to identify factors associated with step numbers after discharge; and to examine the association between functional decline and step numbers after discharge.

Design: Prospective observational cohort study conducted in 2015–2017.

Setting and Participants: Older adults (≥ 70 years of age) acutely hospitalized for at least 48 hours at internal, cardiology, or geriatric wards in 6 Dutch hospitals.

Methods: Steps were counted using the Fitbit Flex accelerometer during hospitalization and 1 week after discharge. Demographic, somatic, physical, and psychosocial factors were assessed during hospitalization. Functional decline was determined 1 month after discharge using the Katz activities of daily living index.

Results: The analytic sample included 188 participants [mean age (standard deviation) 79.1 (6.7)]. One month postdischarge, 33 out of 174 participants (19%) experienced functional decline. The median number of steps was 656 [interquartile range (IQR), 250–1146] at the last day of hospitalization. This increased to 1750 (IQR 675–4114) steps 1 day postdischarge, and to 1997 (IQR 938–4098) steps 7 days postdischarge. Age [$\beta = -57.93$; 95% confidence interval (CI) -111.15 to -4.71], physical performance ($\beta = 224.95$; 95% CI 117.79–332.11), and steps in hospital ($\beta = 0.76$; 95% CI 0.46–1.06) were associated with steps postdischarge. There was a significant association between step numbers after discharge and functional decline 1 month after discharge ($\beta = -1400$; 95% CI -2380 to -420 ; $P = .005$).

Conclusions and Implications: Among acutely hospitalized older adults, step numbers double 1 day postdischarge, indicating that their capacity is underutilized during hospitalization. Physical performance and physical activity during hospitalization are key to increasing the number of steps

This study is funded by the Netherlands Organisation for Health Research and Development (NWO-ZonMw), grant number 16156071.

The organization funding this study had no role in the design or conduct of the study; in the collection, management, analysis, or interpretation of the data; or in the preparation, review, or approval of the manuscript.

The authors declare no conflicts of interest.

* Address correspondence to Daisy Kolk MSc, Amsterdam UMC, University of Amsterdam, Department of Internal Medicine, Section of Geriatric Medicine, D3-335, PO Box 22600, 1100 DD Amsterdam, the Netherlands.

E-mail address: d.kolk@amsterdamumc.nl (D. Kolk).

[†] Contributed equally to this work.

[‡] In addition to the authors, the Hospital-ADL study group consists of the following members: Ingeborg Kuper MD, Annemarieke de Jonghe MD, PhD, Maïke Leguit-Elberse RN, Ad Kamper MD, PhD, Nynke Posthuma MD, PhD, Nienke Brendel MD, and Johan Wold MD.

<https://doi.org/10.1016/j.jamda.2020.06.027>

1525-8610/© 2020 The Authors. Published by Elsevier Inc. on behalf of AMDA – The Society for Post-Acute and Long-Term Care Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

postdischarge. The number of steps 1 week after discharge is a promising indicator of functional decline 1 month after discharge.

© 2020 The Authors. Published by Elsevier Inc. on behalf of AMDA – The Society for Post-Acute and Long-Term Care Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Physical activity is very limited in hospitalized older adults^{1–5} and is associated with functional decline,^{1,6–8} readmissions,^{9,10} and mortality.^{4,11,12} Functional decline, described as new or additional difficulties performing 1 or more activities of daily living (ADL),¹³ is a major problem and is highly prevalent in older adults after acute hospitalization.^{8,14–17} More than 30% of acutely hospitalized older adults experience functional decline at discharge, and only one-third of these adults recover their premorbid functional level 1 month after discharge.¹⁴ The first month after discharge is a critical recovery period, during which new functional disabilities have a high risk of becoming permanent.¹⁴

To prevent functional decline, the World Health Organization recommends keeping older adults as physically active as their abilities and conditions allow.¹⁸ However, physical condition and the ability to be physically active are greatly affected in acutely ill older adults during hospitalization.¹⁹ It is known that physical activity is reduced during hospitalization, but levels of physical activity after discharge have not been well studied.²⁰ Physical activity, defined as “any bodily movement produced by skeletal muscles that requires energy expenditure,”²¹ can be objectively monitored as step numbers in older adults, using accelerometers.^{4,6,10,20} One study using accelerometers showed that taking less than 900 steps per day during hospitalization is associated with functional decline at discharge.⁶ Several studies have shown that physical activity interventions can improve physical performance in older adults,^{15,22–24} but reversing functional decline in the first month after discharge remains challenging.¹⁵

To optimize in-hospital and postdischarge rehabilitation strategies to prevent or reverse eventual functional decline, we need more information on the number of steps taken after discharge and how the number of steps taken relates to functional decline. A previous study²⁰ showed that the number of steps taken in the first week after discharge was associated with 30-day readmission in older adults, indicating that the level of physical activity soon after discharge is a physical marker of readmission risk and overall health. Based on these findings, we hypothesized that step numbers 1 week after discharge are associated with functional decline 1 month after discharge. The aims of this study were to (1) determine step numbers in hospital and up to 1 week after discharge in acutely hospitalized older adults; (2) identify independent predictors of step numbers after discharge; and (3) examine the association between functional decline and step numbers after discharge.

Methods

Study Participants

Participants were from the Hospital-Associated Disability and Impact on Daily Life (Hospital-ADL) Study, a multicenter observational prospective cohort study evaluating the mechanism of hospital-associated functional decline among 401 older adults age 70 years and over, who were acutely admitted to 6 Dutch hospitals for ≥ 48 hours between October 2015 and June 2017.²⁵ Further inclusion criteria were (1) approval of the medical doctor; (2) Mini-Mental State Examination (MMSE) score ≥ 15 ²⁶; and (3) sufficient understanding of the Dutch language to answer the questionnaires. Persons were excluded if they (1) had a life expectancy of less than

3 months or (2) were dependent on help for all 6 basic ADLs (bathing, dressing, eating, toileting, transferring, and maintaining continence).²⁷ There were no further exclusion criteria regarding walking ability. To determine whether a patient was eligible for inclusion, the researcher asked the attending medical doctor for approval before approaching the patient. Participants were recruited from internal medicine, cardiology, and geriatric wards and were asked to provide written consent before inclusion. All participants gave additional consent to wear an activity tracker during hospitalization and after discharge.

The study was approved by the institutional review board. The study was conducted according to the Dutch Medical Research Involving Human Subjects Act and principles of the Declaration of Helsinki (1964). Local approval was provided by all participating hospitals.

Assessments

Demographics, malnutrition, and cognitive functioning of participants were assessed at admission. Comorbidities were retrieved from medical records. Other variables in physical, functional, and psychological domains that might change during hospitalization were assessed at discharge, which was considered the most optimal time point to identify older adults at risk of insufficient recovery postdischarge. Trained researchers administered the standardized study protocol. At 1 month postdischarge, all participants were asked to rate their functionality in performing ADLs.

Counting Steps

The primary outcome was the number of steps taken per day in hospital and up to 1 week after discharge. For the second and third study aims, the primary outcome was the number of steps taken postdischarge. Steps were counted using the Fitbit Flex activity tracker (Fitbit, Inc, San Francisco, CA); an accurate activity tracker to estimate step counts compared with the gold standard Actigraph ($r = 0.96$).²⁸ The Fitbit Flex is worn on the nondominant wrist²⁹ and is user-friendly, which limits study withdrawal.³⁰ The activity tracker was worn continuously from hospital admission to 1 week after discharge. Participants were asked to wear the device at all times, except during charging (1–2 hours per week). Step data were frequently synced to the Fitbit platform and exported at the end of the study. Step numbers were counted every 24 hours, starting at the time of discharge (eg, discharge at 3:00 PM until 3:00 PM the next day was the first day counted after discharge) up to 7 days post discharge. Steps taken in hospital were counted in the same way, backwards from the time of discharge up to the time of study inclusion for a maximum of 7 days. Incomplete days (eg, day of inclusion) and days when no steps were counted were omitted from analyses.

Assessment of Functional Decline

We assessed functional decline based on the ability to perform basic ADLs using the Katz-ADL index score.²⁷ At admission, we asked participants to retrospectively rate their ability to perform ADLs during the 2 weeks before hospital admission. This assessment was repeated at discharge and 1 month after discharge. Participants were

asked whether they needed assistance to perform each ADL. A summary score was calculated ranging from 0 (independent in all ADLs) to 6 (dependent on help for all ADLs). Functional decline was defined as a higher dependency on help in 1 or more ADLs 1 month after discharge compared with baseline (2 weeks before admission).

Assessment of Other Variables

Potential predictors for step numbers were identified based on the Fried theoretical cycle of physical frailty.¹⁹ Comorbidities were assessed with the Charlson comorbidity index (range 0–31), where higher scores indicate higher 1-year mortality risk.³¹ Malnutrition was assessed with the Short Nutritional Assessment Questionnaire and was categorized as no malnutrition, mild malnutrition, and severe malnutrition.³² Handgrip strength was measured 3 times using a dynamometer.^{33,34} The highest score (in kilograms) from both hands was used. Physical performance was assessed with the Short Physical Performance Battery, which measures walking speed, chair stand, and balance. Scores range from 0 to 12, and higher scores indicate better physical performance.³⁵ The number of steps taken in hospital, counted using the Fitbit Flex, was also a potential predictor of steps taken after discharge and was calculated as an average number of steps taken during hospitalization. Functionality in performing ADLs at discharge was assessed using the Katz-ADL, and scores ranged from 0 (independent at all ADLs) to 6 (dependent on all ADLs).²⁷ Somatic geriatric syndromes (fear of falling, pain, fatigue, and mobility impairment), which are prevalent after discharge³⁶ and may impair physical activity, were assessed.³⁷ Fear of falling, pain, and fatigue were measured using the numeric rating scale, which is a continuous scale from 0 (no symptoms) to 10 (severe symptoms).^{38,39} Mobility impairment was assessed using the Functional Ambulation Categories test; scores range from 0 to 5, with higher scores indicating greater dependency.⁴⁰ We assessed psychological geriatric syndromes (cognitive functioning, depressive symptoms, and apathy) because they are (1) highly prevalent in this population³⁶; (2) associated with physical activity⁴¹; and (3) potential barriers for functional recovery after acute hospitalization.^{42,43} Cognitive functioning was assessed with the MMSE, with a score of ≤ 23 indicating cognitive impairment.⁴⁴ Apathy was measured with the Geriatric Depression Scale-3A (GDS-3A) subscale of the GDS-15.⁴⁵ Depressive symptoms were measured with the GDS-15 subscale (GDS-12D).⁴⁶ Higher scores indicated more symptoms of apathy and depression (range, GDS-3A: 0–3; GDS-12D: 0–12).^{45,46}

Statistical Analyses

Statistical analyses were performed in 4 phases: (1) descriptive analysis; (2) linear regression analysis of potential predictors of step numbers post discharge; (3) multivariable regression analysis of potential predictors of step numbers postdischarge using a backwards selection procedure; and (4) multivariable regression analysis of functional decline and step numbers postdischarge. For the first study aim, baseline variables and potential predictors were described with a mean and standard deviation or median and interquartile range (IQR) for continuous variables, and a number (n) and percentage (%) for categorical variables. The primary outcome was number of steps taken per day in hospital and 1 week after discharge, and were presented as medians and IQR. Step numbers were also presented separately for older adults with and without functional decline. We assessed potential predictors of steps numbers taken after discharge using linear mixed models because these can account for correlations between repeated step measures. Factors associated with the primary outcome ($P < .10$) were retained for further analysis. The remaining factors were included in the multivariable linear mixed models. Because of high collinearity between physical factors, we used backward elimination

to identify independent predictors using a cut-off P value of .05.⁴⁷ The results are presented as an unstandardized regression coefficient (beta), 95% confidence interval (CI), and P value. To evaluate the association of functional decline with step numbers, we used linear mixed models and adjusted for the independent predictors of steps numbers. We analyzed functional decline as the independent variable to assess differences in postdischarge step numbers between older adults with and without functional decline.

Mixed linear models can handle missing values in the dependent variable; missing values in the independent variables were imputed.⁴⁸ Based on the missing value patterns and percentage of missing values, we multiply imputed 50 datasets using the multiple imputation chained equations by fully conditional specification with predictive mean matching with K of 10 to the nearest neighbor. Results were pooled using the Rubin rules and used in the linear regression analyses.⁴⁹

Sensitivity analyses were conducted to check for selection bias. All baseline variables were compared between participants included in our analyses vs participants not included in our analyses. In addition, a complete case analysis was performed to identify all risk factors associated with the number of steps taken after discharge. All statistical analyses were performed in Stata SE/15.1 (StataCorp 2017, Stata Statistical Software, Release 15; StataCorp LLC, College Station, TX).

Results

Of the 401 participants, 55 did not consent to wear the activity tracker and postdischarge activity measurements were not available for 158 participants (Figure 1). In total, 188 participants were included in the analysis. The sensitivity analysis showed that participants not included in the analysis ($n = 213$) had a significant lower body mass index, step number during hospitalization, handgrip strength, and the Short Physical Performance Battery score. Participants not included in the analyses also had a longer hospital stay, more frequent cognitive impairment, and a higher Katz-ADL score at discharge.

The mean (standard deviation) age of participants was 79.1 (6.7) years, 106 (56.4%) participants were male, and 169 (89.9%) were born in the Netherlands (Table 1). At discharge, 57 of 188 (30.3%) participants experienced functional decline. Of these, 17 of 48 (35.4%) still had functional decline 1 month after discharge and 9 were lost to follow-up. Of the 131 participants without functional decline at discharge, 16 of 126 (12.7%) had functional decline 1 month after discharge and 5 were lost to follow-up. In total, 33 of 174 (18.9%) of the participants experienced functional decline 1 month postdischarge.

Figure 2, A shows the number of steps per day in hospital and after discharge. The median number of steps was 656 (IQR 250–1146) on the last day before discharge and this more than doubled to a median of 1750 (IQR 675–4114) steps on the first day after discharge. From the second day postdischarge, the number of steps slightly increased to 1997 (IQR 938–4098) on the seventh day postdischarge. Figure 2, B and C show the number of steps separately for older adults with ($n = 33$) and without ($n = 141$) functional decline. In older adults with functional decline, the median was 518 (IQR 229–1541) steps on the last day before discharge, 978 (IQR 437–2395) steps on the first day after discharge, and 965 (IQR 344–2535) steps on the seventh day after discharge. In older adults without functional decline, the median was 1189 (IQR 407–2007) steps on the last day before discharge, 1908 (IQR 763–4421) steps on the first day after discharge, and 2289 (IQR 1222–4727) on the seventh day after discharge.

Table 2 shows the linear regression analysis of the potential predictors and the number of steps after discharge. Age, comorbidities, living independently, cognitive functioning, depressive symptoms, fear of falling, fatigue, handgrip strength, physical performance, malnutrition, functional disabilities, and steps taken in hospital had a P value of $< .10$ and were retained for further analysis. In the




Fig. 1. Derivation of the analytic sample.

multivariable linear regression analysis, we found that age ($\beta = -57.93$; 95% CI -111.15 to -4.71), physical performance ($\beta = 224.95$; 95% CI 117.79 – 332.11), and steps taken in hospital ($\beta = 0.76$; 95% CI 0.46 – 1.06) were independent predictors of physical activity after discharge. This means that for every 1-year increase in age, there was a reduction in steps after discharge, and for every 1-point increase in physical performance and every 1-step increase in hospital, there was an increase in steps taken postdischarge. These results differed slightly to the complete case analysis, which found comorbidities ($\beta = -247.07$; 95% CI -443.84 to -50.3) to also be a significant predictor of steps taken postdischarge.

Table 3 shows the unadjusted and adjusted association of functional decline 1 month after discharge with step numbers after discharge. Older adults with functional decline took significantly fewer steps after discharge than older adults without functional decline did ($\beta = -1400$; 95% CI -2380 to -420 ; $P = .005$). Following adjustment for age, physical performance, and steps taken in hospital, there was no significant association between functional decline and step numbers after discharge ($\beta = -671$; 95% CI -1667 to 325 ; $P = .19$).

Discussion

This study aimed to assess the number of steps taken by older adults in hospital up to 1 week after discharge, to identify factors associated with step numbers after discharge, and to determine if

Table 1

Study Sample Characteristics during Hospitalization (n = 188)

Patient Characteristics	
Age, mean (SD), y	79.1 (6.7)
Male, No. (%)	106 (56.4)
BMI,* mean (SD)	22 (19–25)
Marital status, No. (%)	
Married or living together	106 (56.4)
Widow/widower	27 (14.4)
Single or divorced	55 (29.3)
Born in the Netherlands, No. (%)	169 (89.9)
Education, No. (%)	
Primary school	44 (23.4)
Elementary technical/domestic science school	42 (22.3)
Secondary vocational education	58 (30.9)
Higher level high school/third-level education	44 (23.4)
Polypharmacy, [†] No. (%), (n = 186)	121 (65.1)
Hearing impairment, No. (%)	22 (11.7)
Vision impairment, No. (%)	20 (10.7)
Primary admission diagnosis, No. (%)	
Cardiac	60 (31.9)
Respiratory	31 (16.5)
Other	28 (14.9)
Infection	24 (12.8)
Gastrointestinal	22 (11.7)
Renal	9 (4.8)
Cancer (including hematology)	8 (4.3)
Electrolyte disturbance	6 (3.2)
Length of hospital stay, median (IQR), d	5 (4–8)
Charlson comorbidity index, [‡] median (IQR)	2 (1–3)
SNAQ, No. (%)	
No malnutrition	120 (63.8)
Mild malnutrition	10 (5.3)
Severe malnutrition	58 (30.9)
Handgrip strength, mean (SD), kg, (n = 177)	28.9 (10.6)
Physical performance, [§] median (IQR), (n = 169)	7 (4–10)
Steps taken in hospital, median (IQR), (n = 166)	922 (407–1633)
Katz-ADL score, median (IQR), (n = 182)	1 (0–2)
Fear of falling, median (IQR), NRS score, (n = 183)	0 (0–5)
Pain, median (IQR), NRS score, (n = 184)	0 (0–4)
Fatigue, median (IQR), NRS score, (n = 183)	5 (2–7)
FAC, No. (%), (n = 176)	
Independent	43 (22.9)
Independent on level surfaces	95 (50.5)
Dependent for supervision	28 (14.9)
Dependent for physical assistance I	4 (2.1)
Dependent for physical assistance II	2 (1.1)
Nonfunctional ambulation	4 (2.1)
Cognitive impairment, ^{**} No. (%), (n = 169)	13 (7.7)
Depressive symptoms, ^{††} No. (%), (n = 179)	33 (17.6)
Apathy, ^{‡‡} No. (%), (n = 179)	96 (51.1)
Living independently after discharge, No. (%)	147 (78.2)

BMI, body mass index; FAC, functional ambulation categories; NRS, numeric rating scale; SD, standard deviation; SNAQ, Short Nutritional Assessment Questionnaire.

*Calculated as weight in kilograms divided by height in meters squared.

[†]Use of 5 or more different medications.

[‡]Range of 0–31, with a higher score indicating more or severe comorbidity.

[§]Assessed with the SPPB. The score ranges of 0–12, a higher score indicates a better physical performance.

^{||}Ranging from 0 (independent at all ADLs) to 6 (dependent on all ADLs).

**If a score of <24 on the MMSE.

††If a score of ≥ 6 on the GDS.

‡‡If a score of ≥ 2 on 3 items on the GDS.

functional decline 1 month after discharge is associated with step numbers after discharge. Our results showed that the number of steps taken 1 day after discharge was double those taken prior to discharge. This finding suggests the physical capacity of older adults may be underused during hospitalization, regardless of their (dis)ability. Moreover, the number of steps taken slightly increased for up to 7 days postdischarge, but the number of steps remained low in the majority of older adults. These results demonstrated that, at a younger age, better physical performance and more steps taken in hospital were independently associated with higher postdischarge step




Fig. 2. Number of steps* for (A) all participants (N = 188), (B) participants with functional decline (n = 33), and (C) participants without functional decline (n = 141). p75, the third quartile of the IQR; p25, the first quartile of the IQR. *Day -7 means 7 days prior to discharge; Day 7 means 7 days postdischarge.

Table 2
Univariable (Model A) and Multivariable (Model B) Linear Regression Analyses of Predictors and Step Numbers after Discharge

Variables*	Reference	Model A			Model B				
		β^{\dagger}	95% CI	P Value [‡]	β^{\dagger}	95% CI	P Value [‡]		
Age	Y	−147.68	−200.63	−94.73	<.001	−57.93	−111.15	−4.71	.03
Sex	Male	265.00	−492.3	1023.15	.49				
Charlson comorbidity index	Score	−259.92	−455.90	−63.94	.009				
SNAQ	No malnutrition								
Mild malnutrition		871.23	−770.63	2513.10	.30				
Severe malnutrition		−1292.68	−2097.45	−487.92	.002				
Handgrip strength	kg	59.92	24.88	94.96	.001				
Physical performance	Score	362.57	261.96	463.19	<.001	224.95	117.79	332.11	<.001
Steps taken in hospital	Step	1.07	0.77	1.37	<.001	0.76	0.46	1.06	<.001
Katz-ADL score	Score	−560.73	−853.22	−268.23	<.001				
Fear of falling	Score	−172.00	−293.39	−50.63	.005				
Pain	Score	−64.26	−209.74	81.21	.39				
Fatigue	Score	−184.63	−324.81	−44.46	.01				
FAC	Nonfunctional								
Dependent for physical assistance II		−117.58	−3846.98	3611.82	.95				
Dependent for physical assistance I		365.99	−2686.08	3418.06	.81				
Dependent for supervision		794.6	−1566.12	3155.33	.51				
Independent on level surfaces		1636.18	−621.53	3893.90	.16				
Independent		4215.36	1902.45	6528.27	<.001				
Cognitive impairment	No	−1987.33	−3422.87	−551.78	.007				
Depressive symptoms	No	−1025.03	−2002.02	−48.05	.04				
Apathy	No	−579.28	−1360.52	201.96	.15				
Living independent after discharge	No	1301.93	402.09	2201.77	.005				

FAC, functional ambulation categories; SNAQ, Short Nutritional Assessment Questionnaire.

*See Table 1 for definitions.

[†]Unstandardized regression coefficients for continuous variables are per 1-point increase.

[‡]Variables with a value of < .10 were retained for further analysis.

[§]Using a backward selection procedure variables with a P value of < .05 were included in the model.

numbers. The number of steps taken 1 week after discharge was significantly lower in participants with functional decline 1 month postdischarge compared with in those who experienced no functional decline. This association was not significant after correcting for age, physical performance, and steps taken in hospital.

The median number of steps taken in hospital was consistent with results from previous studies in similar patient populations.^{4,5} We found that approximately 50% of older adults did not take the recommended 900 steps per day during hospitalization, which is associated with a higher risk of functional decline at discharge.⁶ Compared with a study by Fisher et al, we counted fewer steps in the first week postdischarge. A possible explanation for this discrepancy is the different exclusion criteria; in our study, older adults were not excluded if they could not walk safely without assistance, which was an exclusion criterion in the study of Fisher et al.²⁰ Although low step numbers are common during hospitalization in our population,^{4,5} the increase in step numbers that we observed immediately after discharge shows that older adults do not use their full physical capacity during hospitalization. This important finding implies that older adults can be more physically active during hospitalization and underlines the need for in-hospital interventions to stimulate physical activity.^{50,51}

Our findings suggest that, of the of physical frailty factors described by Fried et al,¹⁹ only age,⁴¹ physical performance, and in-hospital step numbers are predictors of postdischarge step numbers. These findings support the theory that acute illness and age-related changes can lead to loss of muscle mass and physical performance, resulting in functional decline and reduced physical activity.¹⁹ In this light, our results suggest that particularly frail older adults, who are more likely to be functionally vulnerable to an acute illness,⁵² become less active after an acute illness.²⁰ Offering targeted interventions to these vulnerable individuals to improve their physical performance and physical activity while they are in hospital may improve postdischarge activity levels.¹⁵

The present study has also shown that a large proportion of acutely hospitalized older adults experience functional decline 1 month after discharge.¹⁴ We also showed that these older adults accrued significantly fewer steps in the first week after discharge compared with older adults without functional decline after discharge. This association between functional decline and physical activity has already been shown during hospitalization.^{6,8} In line with the findings of Fisher et al, our results support the idea that the level of physical activity in the first week after discharge may predict recovery after acute hospitalization.²⁰ However, our results also suggested there is no direct

Table 3
Unadjusted (Model A) and Adjusted (Model B) Association of Functional Decline after Discharge with Step Numbers after Discharge (n = 174)

Variables*	Reference	Model A			Model B				
		β^{\dagger}	95% CI	P Value	β^{\dagger}	95% CI	P Value		
Functional decline at 1-mo	No functional decline	−1400	−2380	−420	.005	−671	−1668	325	.19
Age	Year					−58	−125	9	.09
Physical performance	Score					188	57	318	.005
Steps taken in hospital	Step					0.9	0.6	1.3	<.001

*See Table 1 for definitions.

[†]Unstandardized regression coefficients are per 1-point increase.

association between physical activity and functional decline, and that age and physical performance may confound this relationship.¹⁹ Further research is needed to determine whether modifiable factors like physical activity and physical performance can predict functional decline or recovery after discharge. This information may lead to the development of more effective interventions to prevent or reverse functional decline during the postdischarge recovery period.

Limitations

This study has several potential limitations. First, we aimed to include a cohort of older adults with a wide range of vulnerability. However, we had to exclude participants without postdischarge activity data, which may have resulted in selection bias; this means that participants who stayed in hospital longer, whose physical performance was poor, and who took fewer steps in hospital, and who might have been more vulnerable as a result, were excluded. This may have led to an underestimation of the observed associations and reduced the generalizability of our findings to these older, more vulnerable adults. Second, we assessed physical activity using the Fitbit Flex activity tracker because of its practical applicability, user-friendly wristband, and high reliability in measuring steps. Fitbit Flex wearing was controlled using logbooks and regular checks. However, the Fitbit Flex does not detect nonwear time, so physical activity would have been underestimated if the participant was not wearing the tracker.

A major strength of this study was the continuous assessment of physical activity during hospitalization up to the first week post-discharge in a heterogeneous cohort of acutely hospitalized older adults. A large cohort of older adults with a wide range of diagnoses was included. During hospitalization, we assessed a broad set of potential predictors of physical activity.

Conclusions and Implications

This study shows that step numbers taken by acutely hospitalized older adults double immediately after discharge, indicating that the physical capacity of older persons is underutilized during hospitalization. Acutely hospitalized older adults with a younger age, better physical performance, and higher in-hospital physical activity levels have better postdischarge physical activity levels. Interventions focusing on physical performance and physical activity during hospitalization may optimize postdischarge physical activity, and should continue during the critical postdischarge recovery period. The level of physical activity in the first week after discharge is a promising indicator of functional decline 1 month after discharge. Further research is needed to determine whether physical activity and physical performance can predict recovery after acute hospitalization.

Acknowledgments

We thank Suzanne Schilder MSc, Angelique Heinen MSc, Robin Kwakman MSc, and Jan Jaap Voigt MSc for assistance with data collection.

References

- Brown CJ, Friedkin RJ, Inouye SK. Prevalence and outcomes of low mobility in hospitalized older patients. *J Am Geriatr Soc* 2004;52:1263–1270.
- Fisher SR, Graham JE, Brown CJ, et al. Factors that differentiate level of ambulation in hospitalised older adults. *Age Ageing* 2012;41:107–111.
- Pedersen MM, Bodilsen AC, Petersen J, et al. Twenty-four-hour mobility during acute hospitalization in older medical patients. *J Gerontol Ser A Biol Sci Med Sci* 2013;68:331–337.
- Ostir GV, Berges IM, Kuo YF, et al. Mobility activity and its value as a prognostic indicator of survival in hospitalized older adults. *J Am Geriatr Soc* 2013;61:551–557.
- Sallis R, Roddy-Sturm Y, Chijioke E, et al. Stepping toward discharge: Level of ambulation in hospitalized patients. *J Hosp Med* 2015;10:384–389.
- Agmon M, Zisberg A, Gil E, et al. Association between 900 steps a day and functional decline in older hospitalized patients. *JAMA Intern Med* 2017;177:272–274.
- Hardy SE, Gill TM. Factors associated with recovery of independence among newly disabled older persons. *Arch Intern Med* 2005;165:106–112.
- Zisberg A, Shadmi E, Gur-Yaish N, et al. Hospital-associated functional decline: The role of hospitalization processes beyond individual risk factors. *J Am Geriatr Soc* 2015;63:55–62.
- Chawla H, Bulathsinghala C, Tejada JP, et al. Physical activity as a predictor of thirty-day hospital readmission after a discharge for a clinical exacerbation of chronic obstructive pulmonary disease. *Ann Am Thorac Soc* 2014;11:1203–1209.
- Fisher SR, Graham JE, Ottenbacher KJ, et al. Inpatient walking activity to predict readmission in older adults. *Arch Phys Med Rehabil* 2016;97:S226–S231.
- Lee IM, Shiroma EJ, Kamada M, et al. Association of step volume and intensity with all-cause mortality in older women. *JAMA Intern Med* 2019;179:1105–1112.
- Buchman AS, Yu L, Boyle PA, et al. Total daily physical activity and longevity in old age. *Arch Intern Med* 2012;172:444–446.
- World Health Organization. ICF international classification of functioning, disability and health. Geneva: WHO; 2001.
- Boyd CM, Landefeld CS, Counsell SR, et al. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. *J Am Geriatr Soc* 2008;56:2171–2179.
- Brown CJ, Foley KT, Lowman JD Jr, et al. Comparison of posthospitalization function and community mobility in hospital mobility program and usual care patients: A randomized clinical trial. *JAMA Intern Med* 2016;176:921–927.
- Gill TM, Allore HG, Gahbauer EA, Murphy TE. Change in disability after hospitalization or restricted activity in older persons. *JAMA* 2010;304:1919–1928.
- Buurman BM, Hoogerduijn JG, de Haan RJ, et al. Geriatric conditions in acutely hospitalized older patients: Prevalence and one-year survival and functional decline. *PLoS One* 2011;6:e26951.
- World Health Organization. Global recommendations on physical activity for health 65 years and above. Geneva: WHO; 2011.
- Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: Evidence for a phenotype. *J Gerontol Ser A Biol Sci Med Sci* 2001;56:M146–M156.
- Fisher SR, Kuo YF, Sharma G, et al. Mobility after hospital discharge as a marker for 30-day readmission. *J Gerontol Ser A Biol Sci Med Sci* 2013;68:805–810.
- World Health Organization. Global recommendations on physical activity for health. Geneva: WHO; 2010.
- Martinez-Velilla N, Casas-Herrero A, Zambom-Ferraresi F, et al. Effect of exercise intervention on functional decline in very elderly patients during acute hospitalization: A randomized clinical trial. *JAMA Intern Med* 2019;179:28–36.
- Marzetti E, Calvani R, Tosato M, et al. Physical activity and exercise as countermeasures to physical frailty and sarcopenia. *Aging Clin Exp Res* 2017;29:35–42.
- Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: The LIFE study randomized clinical trial. *JAMA* 2014;311:2387–2396.
- Reichardt LA, Aarden JJ, van Seben R, et al. Unravelling the potential mechanisms behind hospitalization-associated disability in older patients; the Hospital-Associated Disability and impact on daily Life (Hospital-ADL) cohort study protocol. *BMC Geriatr* 2016;16:59.
- McGivney SA, Mulvihill M, Taylor B. Validating the GDS depression screen in the nursing home. *J Am Geriatr Soc* 1994;42:490–492.
- Katz S, Downs TD, Cash HR, Grotz RC. Progress in development of the index of ADL. *Gerontologist* 1970;10:20–30.
- Alharbi M, Bauman A, Neubeck L, Gallagher R. Validation of Fitbit-Flex as a measure of free-living physical activity in a community-based phase III cardiac rehabilitation population. *Eur J Prev Cardiol* 2016;23:1476–1485.
- Bai Y, Welk GJ, Nam YH, et al. Comparison of consumer and research monitors under semistructured settings. *Med Sci Sports Exer* 2016;48:151–158.
- O'Brien T, Troutman-Jordan M, Hathaway D, et al. Acceptability of wristband activity trackers among community dwelling older adults, 36. New York, NY: Geriatr Nurs; 2015. p. S21–S25.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J Chronic Dis* 1987;40:373–383.
- Kruizenga HM, Seidell JC, de Vet HC, et al. Development and validation of a hospital screening tool for malnutrition: The short nutritional assessment questionnaire (SNAQ). *Clin Nutr (Edinburgh, Scotland)* 2005;24:75–82.
- Roberts HC, Denison HJ, Martin HJ, et al. A review of the measurement of grip strength in clinical and epidemiological studies: Towards a standardised approach. *Age Ageing* 2011;40:423–429.
- Trutschnigg B, Kilgour RD, Reinglas J, et al. Precision and reliability of strength (Jamar vs Biodex handgrip) and body composition (dual-energy X-ray absorptiometry vs bioimpedance analysis) measurements in advanced cancer

- patients. *Appl Physiol Nutr Metab = Physiologie appliquee, nutrition, et metabolisme* 2008;33:1232–1239.
35. Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: Association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol* 1994;49:M85–M94.
 36. van Seben R, Reichardt LA, Aarden JJ, et al. The course of geriatric syndromes in acutely hospitalized older adults: The Hospital-ADL Study. *J Am Med Dir Assoc* 2019;20:152–158.e152.
 37. Brown CJ, Williams BR, Woodby LL, et al. Barriers to mobility during hospitalization from the perspectives of older patients and their nurses and physicians. *J Hosp Med* 2007;2:305–313.
 38. Hwang SS, Chang VT, Cogswell J, Kasimis BS. Clinical relevance of fatigue levels in cancer patients at a Veterans Administration Medical Center. *Cancer* 2002;94:2481–2489.
 39. MacCaffery M, Beebe A. *Pain: Clinical manual for nursing practice*. Mosby: St. Louis, Missouri; 1989.
 40. Holden MK, Gill KM, Magliozzi MR, et al. Clinical gait assessment in the neurologically impaired. Reliability and meaningfulness. *Phys Ther* 1984;64:35–40.
 41. Clarke CL, Sniehotta FF, Vadiveloo T, et al. Factors associated with change in objectively measured physical activity in older people: Data from the physical activity cohort Scotland study. *BMC Geriatr* 2017;17:180.
 42. Sands LP, Yaffe K, Covinsky K, et al. Cognitive screening predicts magnitude of functional recovery from admission to 3 months after discharge in hospitalized elders. *J Gerontol Ser A Biol Sci Med Sci* 2003;58:37–45.
 43. Deer RR, Goodlett SM, Fisher SR, et al. A randomized controlled pilot trial of interventions to improve functional recovery after hospitalization in older adults: Feasibility and adherence. *J Gerontol Ser A Biol Sci Med Sci* 2018;73:187–193.
 44. Folstein MF, Folstein SE, McHugh PR. Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–198.
 45. Bertens AS, Moonen JE, de Waal MW, et al. Validity of the three apathy items of the Geriatric Depression Scale (GDS-3A) in measuring apathy in older persons. *Int J Geriatr Psychiatry* 2017;32:421–428.
 46. Eurelings LS, van Dalen JW, Ter Riet G, et al. Apathy and depressive symptoms in older people and incident myocardial infarction, stroke, and mortality: A systematic review and meta-analysis of individual participant data. *Clin Epidemiol* 2018;10:363–379.
 47. Ambler G, Brady AR, Royston P. Simplifying a prognostic model: A simulation study based on clinical data. *Stat Med* 2002;21:3803–3822.
 48. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med* 2011;30:377–399.
 49. White IR, Carlin JB. Bias and efficiency of multiple imputation compared with complete-case analysis for missing covariate values. *Stat Med* 2010;29:2920–2931.
 50. Meesters J, Conijn D, Vermeulen HM, Vliet Vlieland T. Physical activity during hospitalization: Activities and preferences of adults versus older adults. *Physiother Theory Pract* 2019;35:975–985.
 51. Tran K, Wright MD. Senior friendly hospital care: A review of guidelines. In: *CADTH Rapid Response Reports*. Ottawa (ON): Canadian Agency for Drugs and Technologies in Health; 2019.
 52. Fimognari FL, Pierantozzi A, De Alfieri W, et al. The severity of acute illness and functional trajectories in hospitalized older medical patients. *J Gerontol Ser A Biol Sci Med Sci* 2017;72:102–108.