Body weight-supported bedside treadmill training facilitates ambulation in ICU patients

an interventional proof of concept study

Author(s)

Sommers, Juultje; Wieferink, DC; Nollet, F; Dongelmans, Dave A; Engelbert, R.H.H.; van der Schaaf, M.

Publication date

2017

Document Version

Final published version

License

CC BY

[Link to publication](#)

Citation for published version (APA):

Sommers, J., Wieferink, DC., Nollet, F., Dongelmans, D. A., Engelbert, R. H. H., & van der Schaaf, M. (2017). *Body weight-supported bedside treadmill training facilitates ambulation in ICU patients: an interventional proof of concept study*. Poster session presented at WCPT , Capetown.

<http://www.abstractstosubmit.com/wcpt2017/abstracts/main.php?do=YToyOntzOjU6Im1vZHVsljtzOjY6ImRldGFpbCI7czo4OiJkb2N1bWVudC17aTo1MDY7fQ==&>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the library: <https://www.amsterdamuas.com/library/contact/questions>, or send a letter to: University Library (Library of the University of Amsterdam and Amsterdam University of Applied Sciences), Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Body weight-supported bedside treadmill training facilitates ambulation in ICU patients: An interventional proof of concept study

J. Sommers, PT, MSc¹, D.C. Wieferink, PT¹, D.A. Dongelmans, MD, PhD.², F. Nollet, MD, PhD.¹, RHH. Engelbert, PT, PhD.^{1,3}, M. van der Schaaf, PT, PhD.^{1,3}

1: Department of Rehabilitation, 2: Department of Intensive Care Academic Medical Center, University of Amsterdam, The Netherlands.

3: ACHIEVE-Centre of Applied Research, Faculty of Health University of Applied Sciences, Amsterdam, The Netherlands.

Journal of Critical Care, 2017

Background

- Intensive Care Unit-acquired weakness (ICU-AW) is associated with short- and long- term physical impairments and impaired functional status.
- Early mobilisation and ambulation of patients admitted to the ICU improve functional recovery.
- Severe weakness in combination with tubes, lines and machinery are practical barriers for the implementation of ambulation of critically ill patients.

Objective

- To explore the feasibility of Body Weight- Supported Treadmill Training (BWSTT) at the bedside in the ICU.

Methods

- Single centre interventional study
- Tertiary 34-bed mixed medical-surgical ICU of the Academic Medical Center, University of Amsterdam, the Netherlands

Inclusion criteria:

- Adult ICU-patients, mechanically ventilated for ≥ 48 hours
- No contra-indication for mobilisation or activation
- Able to follow instructions
- Muscle strength m. quadriceps $MRC \geq 2$
- Independent sitting balance

Intervention

- The BWST enables ambulation at the bedside within the range of ventilator tubes, lines and monitoring equipment.
- In patients with insufficient motor control or muscle strength for ambulation, a harness is used in combination with a weight bearing construction.

Measurements

- Feasibility was evaluated according to:
 - Eligibility
 - Successful number of BWSTT
 - Number of staff needed
 - Adverse events
 - Number of patients that could not have walked without BWSTT
 - Patient satisfaction and anxiety

Results

Patients characteristics at the first time of BWSTT	
median (IQR) or n	n=20
Age, years	69.5 (52.8 – 77.5)
Male	12 (60%)
Medical category	
• Medical	9 (45%)
• Non-elective surgery	4 (20%)
• Elective surgery	7 (35%)
APACHE II	18 (15 – 20)
Time in ICU to first BWSTT, days	23.0 (10.0 – 56.3)
Mechanical ventilation, days	10.0 (7.1 – 31.5)
MRC sum score	40 (32.5 – 47.5)
ICU-AW (MRC < 48)	15 (75%)
Functional Ambulation Categories (FAC range 0-5)	0.0 (0.0 – 1.0)
Functional Ambulation Categories 0	13 (65%)
Characteristics of ICU treatment devices during BWSTT	
Continuous monitoring	20 (100%)
Patients with mechanical ventilation	4 (20%)
Patients with tracheostomy	8 (40%)
Surgical wounds	11(55%)
-Sternotomy	6 (55%)
-Abdominal	3 (27%)
-neck	2 (18%)
Patients with infusion lines or drains	20 (100%)

Results

- BWSTT was performed in:
 - 20 patients and 54 sessions
 - 53 sessions were successfully performed
- Two numbers off staff needed
- BWSTT is feasible and safe:
 - Median treatment time 25 minutes
 - No adverse events
 - Patients are not anxious
 - Patients are very satisfied
 - All patients were not able to walk or would have walked shorter distance without the BWST

Conclusion

- BWSTT is feasible and safe and facilitates early ambulation with critically ill patients in the ICU.
- To perform BWSTT less staff is needed compared to ambulation without BWSTT.

References

- Hermans G, Crit. Care 2015;19:274.
- Needham DM, Crit Care Med. 2009;37(10 Suppl):S436-S441
- Harrold ME, Crit. Care. 2015;19:336.
- Burtin C, Crit Care Med 2009; 37: 2499-505.
- Sommers J, Clinical Rehabilitation. 2015; DOI: 10.1177/0269215514567156

Contact

Juultje Sommers: j.sommers@amc.uva.nl