

## Amsterdam University of Applied Sciences

### Digitally supported dietary counseling increases protein intake in community dwelling older adults: preliminary results of the VITAMIN RCT

van den Helder, J.; van Dronkelaar, C.; Tieland, M.; Weijs, P.J.M.

**Publication date**

2018

**Document Version**

Final published version

[Link to publication](#)

**Citation for published version (APA):**

van den Helder, J., van Dronkelaar, C., Tieland, M., & Weijs, P. J. M. (2018). *Digitally supported dietary counseling increases protein intake in community dwelling older adults: preliminary results of the VITAMIN RCT*. Poster session presented at Science Exchange Day, Amsterdam, Netherlands.

**General rights**

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

**Disclaimer/Complaints regulations**

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the library:

<https://www.amsterdamuas.com/library/contact/questions>, or send a letter to: University Library (Library of the University of Amsterdam and Amsterdam University of Applied Sciences), Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

# DIGITALLY SUPPORTED DIETARY COUNSELING INCREASES PROTEIN INTAKE IN COMMUNITY DWELLING OLDER ADULTS

J. van den Helder<sup>1,2</sup>, C. van Dronkelaar<sup>1</sup>, M. Tieland<sup>1</sup>, P.J.M. Weijs<sup>1,3</sup> and VITAMIN research group<sup>1,2,3,4</sup>

<sup>1</sup> Faculty of Sports and Nutrition, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands; <sup>2</sup> Amsterdam Center for Innovative Health Practice (ACHIEVE), Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands; <sup>3</sup> Department of Nutrition and Dietetics, Amsterdam University Medical Centers, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands; <sup>4</sup> CREATE-IT Applied Research, Faculty of Digital Media and Creative Industries, Amsterdam University of Applied Sciences, Amsterdam, The Netherlands.

## Rationale

In order to prevent sarcopenia in community dwelling older adults a higher daily protein intake is needed. A new e-health strategy for dietary counseling was used with the aim to increase total daily protein intake to optimal levels (minimal 1.2 g/kg/day, optimal 1.5 g/kg/day) through use of regular food products.


## Methods

The VITAMIN (VITal AMsterdam older adults IN the city) RCT included 245 community dwelling older adults (age ≥ 55y): control, exercise, and exercise plus dietary counseling (protein) group. Dietary intake was measured by a 3-day dietary record at baseline and after 6 months intervention. In total 173 subjects were eligible for analysis. A two-way mixed ANOVA with time, group, and time\*group interaction was performed. Post-hoc Bonferroni was performed with significance level at  $p < 0.05$ .



## Conclusion


This study shows digitally supported dietary counseling improves protein intake sufficiently in community dwelling older adults. Protein intake increase by counseling with e-health is a promising strategy for dietitians and health care professionals in order to support healthy ageing.



**Figure 1. Mean protein intake of the study groups of community dwelling older adults.**

## Results

Mean age of the subjects was  $72.1 \pm 6.3$ , with a BMI of  $25.7 \pm 4.2$  of which 68% were females. ANOVA revealed significant effect of time, group and time\*group ( $p < 0.001$ ). Figure 1 shows higher protein intake over time in the dietary counseling group than either control ( $p = 0.038$ ) or exercise ( $p = 0.008$ ) group. Additional analyses revealed no change in vegetable protein intake. The higher protein intake was fully accounted for by animal protein intake. In the dietary counseling group 72% of subjects increased protein intake above the minimum intake level and 41% of the subjects above optimal level (see Figure 2).



**Figure 2. Achievement of protein intake recommendations for the study groups of community dwelling older adults.**

## Correspondence

[j.e.m.van.den.helder@hva.nl](mailto:j.e.m.van.den.helder@hva.nl) / Jantine van den Helder, Faculty of Sports and Nutrition, Amsterdam University of Applied Sciences, Dokter Meurerlaan 8, 1067 SM Amsterdam, The Netherlands