

The pre-and postoperative course of functional status of patients with esophageal cancer undergoing esophagectomy

a prospective longitudinal study

Author(s)

van Egmond, M.A.; van der Schaaf, M.; Klinkerbijl, J.H.G.; Twisk, J.W.R.; Engelbert, R.H.H.; van Berge Henegouwen, M.I.

Publication date

2018

[Link to publication](#)

Citation for published version (APA):

van Egmond, M. A., van der Schaaf, M., Klinkerbijl, J. H. G., Twisk, J. W. R., Engelbert, R. H. H., & van Berge Henegouwen, M. I. (2018).

The pre-and postoperative course of functional status of patients with esophageal cancer undergoing esophagectomy: a prospective longitudinal study. Poster session presented at Dag van de fysiotherapeut 2018.



General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the library: <https://www.amsterdamuas.com/library/contact/questions>, or send a letter to: University Library (Library of the University of Amsterdam and Amsterdam University of Applied Sciences), Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

The pre- and postoperative course of functional status of patients with esophageal cancer undergoing esophagectomy

A prospective longitudinal study

M.A. van Egmond^{1,2,3}, M. van der Schaaf^{1,2}, J.H.G. Klinkenbijn^{5,6}, J.W.R. Twisk⁷, R.H.H. Engelbert^{1,2}, M.I. van Berge Henegouwen⁴

1 Amsterdam UMC, University of Amsterdam, Department of Rehabilitation, Amsterdam Movement Sciences, the Netherlands; 2 ACHIEVE - Centre of Applied Research, Faculty of Health, Amsterdam University of Applied Sciences, Amsterdam, the Netherlands; 3 European School of Physiotherapy, Amsterdam University of Applied Sciences, Faculty of Health, Amsterdam, The Netherlands; 4 Amsterdam UMC, Department of Surgery, University of Amsterdam, Amsterdam, The Netherlands; 5 Gelre Hospital Apeldoorn-Zutphen, Department of Surgery, The Netherlands; 6 University of Amsterdam, Amsterdam, The Netherlands; 7 Department of Epidemiology and Biostatistics, Amsterdam University Medical Centers, Amsterdam, The Netherlands

Introduction

- The prevalence of postoperative complications (POC) in patients with esophageal cancer undergoing esophagectomy is high (up to 60%).¹
- Physiotherapy is recommended to improve functional status in the pre- and postoperative stages. However, knowledge of the course of functional status and its association with POC in patients after esophagectomy is limited.²

Objective

To investigate the course of functional status in patients with esophageal cancer before and after esophagectomy.

Methods

Patients: We assessed 155 (100%) patients at 3 months and at 1 day before surgery, 109 (70.3%) at 1 week after surgery and 60 (55%) at 3 months after surgery (Figure 1). We used linear mixed model analyses.




Figure 1. Measurements over time

Functional status: was assessed according to the domains of the International Classification of Functioning, Disability and Health (ICF) (Figure 2).




Figure 2. Measurements of functional status

Results

- The incidence of POC was 83 (53.5%).
- The mean level of HGS, proximal muscle strength of the lower extremities and MIP at T0 and T1 was higher than predicted when adjusted for sex and age (Table 1).
- At T4, all mean functional status measures returned to baseline levels, but patients had significantly less HGS and experienced more fatigue (Table 1).
- No differences in functional status between patients with and without complications at the different time points were observed, apart from 2 MWT at T2.

Measurements	T1	T2	Δ T2-T1 (95% CI)	T3	Δ T3-T1 (95% CI)	T4 (95% CI)	Δ T4-T1 (95% CI)
HGS	110.99 (25.88)	114.26 (21.67)	2.61 (-1.29, 6.51)	108.17 (23.32)	-2.75 (-6.83, 1.32)	104.81 (30.05)	-6.16 (-11.25, -1.06)
MIP	119.74 (40.63)	125.36 (38.49)	4.06 (-6.2, 8.74)	88.42 (33.56)	-37.21 (-43.04, -31.38)	120.41 (38.11)	-4.38 (-10.52, 1.77)
30 s CST	118.08 (28.85)	117.00 (38.16)	-2.78 (-4.69, 5.25)	87.64 (34.00)	-32.23 (-37.73, -26.73)	119.13 (38.07)	-2.33 (-8.87, 4.21)
2 MWT (m)	193.55 (29.22)	202.26 (35.56)	9.36 (4.04, 14.68)	147.52 (37.17)	-48.35 (-54.00, -42.70)	191.20 (34.42)	-5.72 (-12.63, 1.19)
MFI Fatigue, median (IQR)	45.45 (26.25)	40.84 (23.75)	-4.65 (-7.58, -1.72)	-	-	49.20 (17.14)	4.69 (7.1, 8.67)
EORTC QLQ C30 Quality of Life (% median (IQR))	73.44 (15.73)	80.53 (14.16)	7.10 (3.92, 10.27)	-	-	71.81 (18.11)	-2.46 (-6.69, 1.76)
LAPAQ (Kcal/day) median (IQR)	475.37 (550.23)	864.48 (671.00)	363.96 (177.58, 550.33)	-	-	453.04 (539.36)	136.84 (-106.82, 380.50)

Table 1. Differences in functional status outcomes

Conclusion

This study shows that the functional status of high-risk surgical patients undergoing esophagectomy returned to baseline values three months after surgery, despite the high incidence of postoperative complications.

Practical implications

This study illustrates the importance of first evaluating the pre- and postoperative course of the functional status of high-risk surgical patients and of timing and tailoring the physiotherapy treatment to a patient's individual needs, before referring them for pre- or postoperative physiotherapy.

References

- Feeney C, Reynolds JV, Hussey J. Preoperative physical activity levels and postoperative pulmonary complications post-esophagectomy. *Dis Esophagus* 2011 Sep;24(7):489-494.
- Hoogeboom TJ, Dronkers JJ, Hulzebos EH, van Meeteren NL. Merits of exercise therapy before and after major surgery. *Curr Opin Anaesthesiol* 2014 Apr;27(2):161-166.