

Amsterdam University of Applied Sciences

Comprehensive geriatric assessment and transitional care in acutely hospitalized patients

the transitional care bridge randomized clinical trial

Buurman, Bianca M.; Parlevliet, Juliette L.; Allore, Heather G.; Blok, Willem; van Deelen, Bob A.J.; Moll van Charante, Eric P.; de Haan, Rob J.; de Rooij, Sophia E.

DOI

[10.1001/jamainternmed.2015.8042](https://doi.org/10.1001/jamainternmed.2015.8042)

Publication date

2016

Document Version

Final published version

Published in

JAMA internal medicine

[Link to publication](#)

Citation for published version (APA):

Buurman, B. M., Parlevliet, J. L., Allore, H. G., Blok, W., van Deelen, B. A. J., Moll van Charante, E. P., de Haan, R. J., & de Rooij, S. E. (2016). Comprehensive geriatric assessment and transitional care in acutely hospitalized patients: the transitional care bridge randomized clinical trial. *JAMA internal medicine*, 176(3), 302-309.
<https://doi.org/10.1001/jamainternmed.2015.8042>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Library know, stating your reasons. In case of a legitimate complaint, the Library will make the material inaccessible and/or remove it from the website. Please contact the library:

<https://www.amsterdamuas.com/library/contact/questions>, or send a letter to: University Library (Library of the University of Amsterdam and Amsterdam University of Applied Sciences), Secretariat, Singel 425, 1012 WP Amsterdam, The Netherlands. You will be contacted as soon as possible.

Original Investigation

Comprehensive Geriatric Assessment and Transitional Care in Acutely Hospitalized Patients

The Transitional Care Bridge Randomized Clinical Trial

Bianca M. Buurman, RN, PhD; Juliette L. Parlevliet, MD; Heather G. Allore, PhD; Willem Blok, MD, PhD; Bob A. J. van Deelen, MD; Eric P. Moll van Charante, MD, PhD; Rob J. de Haan, RN, PhD; Sophia E. de Rooij, MD, PhD

[+ Supplemental content at
jamainternalmedicine.com](#)

IMPORTANCE Older adults acutely hospitalized are at risk of disability. Trials on comprehensive geriatric assessment (CGA) and transitional care present inconsistent results.

OBJECTIVE To test whether an intervention of systematic CGA, followed by the transitional care bridge program, improved activities of daily living (ADLs) compared with systematic CGA alone.

DESIGN, SETTING, AND PARTICIPANTS This study was a double-blind, multicenter, randomized clinical trial conducted at 3 hospitals with affiliated home care organizations in the Netherlands between September 1, 2010, and March 1, 2014. In total, 1070 consecutive patients were eligible, 674 (63.0%) of whom enrolled. They were 65 years or older, acutely hospitalized to a medical ward for at least 48 hours with an Identification of Seniors at Risk–Hospitalized Patients score of 2 or higher, and randomized using permuted blocks stratified by study site and Mini-Mental State Examination score (<24 vs ≥24). The dates of the analysis were June 1, 2014, to November 15, 2014.

INTERVENTIONS The transitional care bridge program intervention was started during hospitalization by a visit from a community care registered nurse (CCRN) and continued after discharge with home visits at 2 days and at 2, 6, 12, and 24 weeks. The CCRNs applied the CGA care and treatment plan.

MAIN OUTCOMES AND MEASURES The main outcome was the Katz Index of ADL at 6 months compared with 2 weeks before admission. Secondary outcomes were mortality, cognitive functioning, time to hospital readmission, and the time to discharge from a nursing home.

RESULTS The study cohort comprised 674 participants. Their mean age was 80 years, 42.1% (n = 284) were male, and 39.2% (n = 264) were cognitively impaired at admission. Intent-to-treat analysis found no differences in the mean Katz Index of ADL at 6 months between the intervention arm (mean, 2.0; 95% CI, 1.8-2.2) and the CGA-only arm (mean, 1.9; 95% CI, 1.7-2.2). For secondary outcomes, there were 85 deaths (25.2%) in the intervention arm and 104 deaths (30.9%) in the CGA-only arm, resulting in a lower risk on the time to death within 6 months after hospital admission (hazard ratio, 0.75; 95% CI, 0.56-0.99; P = .045; number needed to treat to prevent 1 death, 16). No other secondary outcome was significant.

CONCLUSIONS AND RELEVANCE A systematic CGA, followed by the transitional care bridge program, showed no effect on ADL functioning in acutely hospitalized older patients.

TRIAL REGISTRATION Netherlands Trial Registry: [NTR2384](#)

JAMA Intern Med. 2016;176(3):302-309. doi:10.1001/jamainternmed.2015.8042
Published online February 15, 2016.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Bianca M. Buurman, RN, PhD, Section of Geriatric Medicine, Department of Internal Medicine, Academic Medical Center, University of Amsterdam, PO Box 22660, 1100 DD Amsterdam, the Netherlands
(b.m.vanes@amc.nl).

Within 6 months of acute hospitalization, 30% to 50% of older patients experience a loss of essential activities of daily living (ADLs),¹⁻³ while 20% to 30% are readmitted⁴ and 20% to 30% die.^{1,5} Two clinical care models specifically target older persons at risk for these negative outcomes during hospitalization and in the transition from hospital to home, including the comprehensive geriatric assessment (CGA) team approach⁶ and transitional care.⁷

In the CGA team approach, a multidisciplinary team visits older patients during hospitalization, performs a CGA focused on illnesses and geriatric conditions, initiates interventions, and monitors each patient until hospital discharge. The effectiveness of this approach has been studied extensively, with one meta-analysis⁶ demonstrating a positive effect on cognitive functioning, while another meta-analysis⁸ found a reduction in mortality until 8 months after discharge. On most other outcomes, the CGA team approach has not demonstrated effectiveness.^{6,8} These mixed findings may be due to nonadherence to recommendations provided in the CGA care and treatment plans, and the care initiated during hospitalization may be discontinued on discharge regardless of the presence or worsening of the geriatric conditions.

Transitional care aims to ensure safe transitions between care settings.⁷ In the transition from hospital to home, many patients experience adverse drug events,⁹ have inadequate follow-up,¹⁰ and manifest difficulties with the execution of discharge instructions.¹¹ Transitional care is a time-limited service. After a number of visits, this care is discontinued, and the patient is handed over to primary care. Transitional care has shown beneficial effects on readmission rates.^{12,13} Nurse care coordination, a home visit within 2 days after hospital discharge, and communication between the hospital and primary care providers are intervention components associated with a reduction in readmission.¹² However, the effects of transitional care on mortality are inconsistent,¹⁴⁻¹⁶ and only one study¹⁴ to date has studied the effect on ADLs.

Combining the 2 clinical care models might be beneficial for older patients by assisting during hospitalization, by aiding in their transition home, and by providing home follow-up for geriatric conditions after hospital discharge. In this double-blind, multicenter, randomized clinical trial, we tested whether an intervention of an in-hospital systematic CGA by a geriatric consultation team, followed by a transitional care program, reduced ADL disabilities by 6 months after discharge in older persons who were acutely admitted to a medical ward compared with older persons who received only systematic CGA during hospitalization.

Methods

Design and Setting

Three hospitals with affiliated home care organizations in the Netherlands participated in the transitional care bridge program, a multicenter, double-blind (via the postponed informed consent procedure),¹⁷ randomized clinical trial conducted between September 1, 2010, and March 1, 2014. The participating hospitals in the Netherlands included the Aca-

demical Medical Center in Amsterdam (a 1024-bed university teaching hospital), the Onze Lieve Vrouwe Gasthuis in Amsterdam (a 555-bed teaching hospital), and the Flevo Hospital in Almere (a 386-bed regional teaching hospital). All these hospitals had a geriatric consultation team. A community care registered nurse (CCRN) from Cordaan Home Care, Buurtzorg Nederland, and Zorggroep Almere provided the transitional care bridge program to participants who were randomized to the intervention group. The trial protocol has been previously described¹⁸ and can be found in [Supplement 1](#). The eMethods 1 in [Supplement 2](#) describes the health care system in the Netherlands.

Ethics

The institutional review board of the Academic Medical Center approved this study (protocol ID MEC10/O82). The institutional review boards of the Onze Lieve Vrouwe Gasthuis and Flevo Hospital provided local approval.

Participants

Consecutive patients who were 65 years or older, were acutely admitted for at least 48 hours to an internal medicine department, and were at risk for functional decline were eligible for inclusion in this study. The risk of functional decline was assessed using the Identification of Seniors at Risk-Hospitalized Patients (ISAR-HP),¹⁹ and those with a score of 2 or higher were considered to be at increased risk for functional decline (eMethods 2 in [Supplement 2](#)).

Recruitment of Participants and Informed Consent

Experienced trained research nurses met with all consecutive patients within 48 hours of admission. Eligible patients were invited to participate in the trial and provided written informed consent using a postponed informed consent procedure.¹⁷ For those patients with severe cognitive impairment due to dementia or delirium (Mini-Mental State Examination [MMSE] score, <16)²⁰ or severe acute illness (eg, shock), their health care proxy provided written informed consent.

Baseline Data Collection

The baseline assessment included the following: demographics, the ISAR-HP, premorbid ADL functioning 2 weeks before admission (6-item original Katz Index of ADL²¹), cognitive functioning (MMSE score²⁰), geriatric conditions (eg, polypharmacy, incontinence, malnutrition,²² delirium,²³ and fall risk), all admission diagnoses, and comorbidity measured at discharge (Charlson Comorbidity Index²⁴). A complete overview of the instruments used is described elsewhere.¹⁸

Randomization

Consented participants were randomly assigned to the intervention arm (systematic CGA and the transitional care bridge program) or the control arm (systematic CGA alone). The randomization list was created using an online system (TENALEA Clinical Trial Data Management System; Trans European Network for Clinical Trial Services), with a maximum permuted block size of 20 and stratification by study site and MMSE score (<24 vs ≥24). To ensure allocation concealment,

Table 1. Components of the Transitional Care Bridge Program Intervention Arm and the Systematic CGA-Only Control Arm

Time Frame	Intervention Component	Professional Involved	Intervention	CGA Only
≤48 h After hospital admission	CGA ^a	Geriatric-trained RN with geriatrician	X	X
	Care and treatment plan, with prioritization by patient and geriatric team	Geriatric consultation team	X	X
During hospital stay	Multidisciplinary care provided by geriatric team	RN and geriatrician; as-needed physical therapist, dietician, and team on the ward	X	X
Before hospital discharge	In-person handover of CGA and care and treatment plan by geriatric team to CCRN	Geriatric consultation team, team on the ward, CCRN	X	
Before hospital discharge	Visit of CCRN to participant	CCRN	X	
≤2 d After hospital discharge	Home visit or visit to the NH, with medication reconciliation; initiate home care; check for activities of daily living and availability of adequate help; target geriatric conditions; contact GP or NH physician	CCRN	X	
2 wk	Home visit or visit to the NH, evaluation of hospital-based care and treatment plan, goal setting by participant, protocol adherence for incident and prevalent geriatric conditions	CCRN	X	
6 wk	Home visit or visit to the NH, focusing on the proxy and caregiver burden; further actions on geriatric conditions	CCRN	X	
12 wk	Home visit or visit to the NH, focusing on care needs for incident and prevalent geriatric conditions	CCRN	X	
24 wk	Home visit or visit to the NH, focusing on care needs; evaluation of geriatric conditions; handover to GP	CCRN	X	

Abbreviations: CCRN, community care registered nurse; CGA, comprehensive geriatric assessment; GP, general practitioner; NH, nursing home; RN, registered nurse.

^a The CGA was conducted by a geriatric-trained RN, who assessed participants on 18 geriatric conditions using a standardized geriatric assessment form. After this, participants were asked whether they recognized the detected geriatric conditions, if they needed any (additional) care or treatment, and which conditions should have priority in the care and treatment plan. The geriatric RN presented these findings to the team's geriatrician. If a psychological problem was present (eg, delirium, cognitive impairment [Mini-Mental State Examination score <24]), or depression) or if the participant had 5 or more geriatric conditions, the geriatrician visited the

participant. A personalized care and treatment plan was initiated, and participants were followed up throughout hospitalization. The geriatric consultation team consisted of at least a geriatrician, geriatric RN, physical therapist, and dietician. After discharge, the medical responsibility was handed over to the GP. In the intervention group, the CCRN contacted the GP in case of problems with medications or geriatric conditions. There was no predefined contact moment between the CCRN and GP. All GPs of patients in the intervention arm received the CGA results and a letter informing them about the transitional care bridge, including the name and telephone number of the CCRN who was coordinating the transitional care bridge program for the patient.

the research nurse used a secure randomization website to receive the treatment allocation when submitting patient information.

Treatment Arms

Components of the systematic CGA and the transitional care bridge program are listed in Table 1. The CCRNs who conducted the transitional care bridge program received additional training before the start of the intervention (eMethods 3 in Supplement 2). All randomized participants received a systematic CGA within 48 hours of admission (Table 1).

Transitional Care Bridge Program Intervention

After the geriatric-trained registered nurse conducted the CGA, the CCRN was contacted to visit the hospital to receive a personal handover of the CGA, to initiate the personalized care and treatment plan (CTP), and to meet with the participant and informal caregiver to discuss their needs. This visit was performed as soon as possible, occurring a median of 4 days after admission.

After discharge, the CCRN at a home visit 2 days later performed medication reconciliation, answered the participant's questions, and completed a needs assessment. If a participant was discharged to a nursing home, the CCRN also visited the nursing home within 2 days after discharge.

At weeks 2, 6, 12, and 24 after discharge, the CCRN performed home visits or visits to the nursing home, with the CTP providing the guidance for these home visits. Geriatric conditions were monitored, and interventions were continued or newly initiated. For geriatric conditions, evidence-based intervention protocols were provided (eMethods 4 in Supplement 2).

Outcomes

The original Katz Index of ADL²¹ at 6 months was the primary end point. The participants were asked whether they needed help to perform each ADL activity 2 weeks before admission and at the time of the follow-up assessments. The Katz Index of ADL ranges from 0 to 6, with higher scores indicating more dependence. Secondary outcomes included mortality at 1 month and 6 months after admission. Complete data on dates of death were verified using electronic medical records or the municipal registry. Cognitive functioning was measured at baseline and at 6 months with the MMSE. Other secondary outcomes were the time to the first unplanned hospital readmission (within 6 months) and, among those discharged to a nursing home, the time to discharge from the nursing home to the community.

Data Collection and Masking

The geriatric-trained registered nurse performed the screening and baseline data collection, which was conducted before the randomization because all participants received CGA. Re-

search assistants masked to the treatment allocation conducted all outcome assessments, including an in-home assessment at 6 months. If a participant was still in a nursing home, the assessment was performed there.

Adherence to the Intervention Protocol

The CCRNs registered each visit for the participants in the intervention arm. Based on these logs and predefined quality indicators (eMethods 5 in Supplement 2), we calculated adherence to the intervention protocol (eTable 1 in Supplement 2).

Sample Size Calculation

The trial was powered for the primary end point of the difference in the Katz Index of ADL between the control and intervention arms at 6 months compared with that at 2 weeks before hospital admission, assuming an effect size of 0.25 to represent a clinically important change of 0.5 point on the original Katz Index of ADL. In total, 256 participants were needed for each treatment arm to achieve 80% power with a 2-sided type I error of 5% and standard deviation of the Katz Index of ADL change of 2.0. The expected attrition due to mortality was 25%. Therefore, 674 participants were enrolled.

Statistical Analysis


All primary analyses were performed on an intent-to-treat basis. Descriptive characteristics of each arm were calculated using proportions or means and SDs, as appropriate. The ADLs 2 weeks before baseline were compared between survivors and decedents with Wilcoxon median tests. All outcome models were adjusted for the randomization strata of study site and the MMSE score (<24 vs ≥24). The primary outcome of ADL functioning was analyzed using a linear mixed-effects model with participant-specific random intercepts, time (baseline or 6 months), treatment arm, and their interaction. Sensitivity analyses using the above approach were performed to assess the number of ADL disabilities, for which decedents were assigned a disability score of 7.

For the secondary outcome of mortality from the time of hospital admission to 1 month and 5 months was analyzed by Cox proportional hazards regression analysis. The number needed to treat was estimated.²⁵ Sensitivity analyses were performed for those who survived past hospital day 4 (the median day when the CCRN started in-hospital visits) to 6 months. For the secondary outcomes of predefined subgroups based on the ISAR-HP risk assessment score (2-3 vs 4-5) and Charlson Comorbidity Index (0-3 vs ≥4), a Cox proportional hazards regression for the time to death tested whether the intervention effect differed by the respective subgroups.

Next, the time to unplanned readmission within 6 months and the time to discharge home (among those discharged to a nursing home) were analyzed using the competing risk models by Fine and Gray,²⁶ accounting for death. Last, cognitive functioning was analyzed by a linear mixed-effects model as described above. All secondary outcomes were adjusted for multiplicity using the method by Hochberg.²⁷

All analyses were performed using statistical software (SAS, version 9.4; SAS Institute Inc). Two-tailed $P < .05$ denoted statistical significance.

Figure 1. CONSORT Diagram of Participant Inclusion



For ADL and cognition, all participants had baseline data and were included in the analyses of these outcomes. The numbers specified are those with 6-month assessment data on ADL and cognition. Nursing home discharged to community applies only for those who went directly from the hospital to a nursing home. Readmission data were only available for those who were discharged alive. ADL indicates activities of daily living; CCRN, community care registered nurse; CGA, comprehensive geriatric assessment; and CONSORT, Consolidated Standards of Reporting Trials.

Results

Patient Inclusion

From October 1, 2010, to January 31, 2013, a total of 1070 consecutive patients were determined to be eligible for the study, and 674 (63.0%) were enrolled (Figure 1). The health care proxies provided informed consent for 55 participants (16.3%) in the intervention arm and for 58 participants (17.2%) in the CGA-only arm. The 2 arms were well balanced on baseline characteristics (Table 2). Overall, the participants had a mean age of 80 years and a mean of 1.8 preexisting ADL disabilities. eTable 2 in Supplement 2 lists additional information on the baseline characteristics for both arms. The median (interquartile range [IQR]) lengths of stay were 8 days (IQR, 5-12 days) in the intervention arm and 8 days (IQR, 5-14 days) in the CGA-only arm. There were 21 hospital deaths (6.2%) in the intervention arm and 34 hospital deaths (10.1%) in the control arm.

Adherence to the Intervention Protocol

Adherence to the intervention protocol is summarized in eTable 1 in Supplement 2. The CGA was conducted for all participants, and a CTP was provided to 95.0% (320 of 337)

Table 2. Baseline Characteristics of the Study Population^a

Variable	Intervention (n = 337)	CGA Only (n = 337)
Study site, No. (%)		
Academic Medical Center	173 (51.3)	173 (51.3)
Onze Lieve Vrouwe Gasthuis	69 (20.5)	70 (20.8)
Flevo Hospital	95 (28.2)	94 (27.9)
Demographics		
Age, mean (SD), y	79.7 (7.3)	80.0 (7.8)
Male sex, No. (%)	142 (42.1)	142 (42.1)
Living situation before admission, No./total No. (%)		
Independent	228/336 (67.9)	224/334 (67.1)
Senior residence	68/336 (20.2)	74/334 (22.2)
Assisted living	26/336 (7.7)	31/334 (9.3)
Other	14/336 (4.2)	5/334 (1.5)
Married, No./total No. (%)	128/334 (38.3)	96/335 (28.7)
Born in the Netherlands, No. (%)	286 (84.9)	289 (85.8)
Health care proxy provided informed consent, No. (%)	55 (16.3)	58 (17.2)
ISAR-HP score, mean (SD) ^b	3.8 (0.9)	3.8 (1.0)
Katz Index of ADL baseline, mean (SD) ^c	1.7 (1.8)	1.8 (1.7)
Charlson Comorbidity Index, mean (SD) ^d	3.11 (2.17)	3.36 (2.40)
Admission diagnosis, No./total No. (%)		
Infection	102 (30.3)	94 (27.9)
Gastrointestinal	49 (14.5)	47 (13.9)
Cardiac	31 (9.2)	41 (12.2)
Respiratory	38 (11.3)	40 (11.9)
Cancer, including hematology	9 (2.7)	22 (6.5)
Electrolyte disturbances	23 (6.8)	16 (4.7)
Renal	11 (3.3)	12 (3.6)
Other ^e	74 (22.0)	65 (19.3)
CGA at Admission, No./Total No. (%)		
Somatic domain		
Polypharmacy ^f	222/277 (80.1)	212/269 (78.8)
Incontinence ^g	96/277 (34.7)	99/279 (35.5)
Indwelling urinary catheter ^h	69/279 (24.7)	68/279 (24.4)
Dizziness ⁱ	129/283 (45.6)	137/281 (48.8)
Malnutrition ^j	179/336 (53.3)	172/337 (51.1)
Pain ^k	175/336 (52.1)	146/316 (46.2)
Psychological domain		
Cognitively impaired ^l	136/337 (40.4)	128/337 (38.0)
Sad or depressive symptoms in the past month ^m	96/279 (34.3)	106/280 (37.9)
Less interest in activities ^m	88/279 (31.5)	110/280 (39.2)
Functional domain		
Fall in the past 6 mo ⁿ	149/329 (45.3)	174/332 (52.4)
Sleep disturbances ^o	164/283 (58.0)	143/280 (51.1)
Hearing impairment ^p	66/261 (25.3)	65/267 (24.3)
Visual impairment ^q	53/270 (19.6)	68/271 (25.1)

Abbreviations: ADL, activities of daily living; CGA, comprehensive geriatric assessment; ISAR-HP, Identification of Seniors at Risk-Hospitalized Patients.

^a Some denominators vary because of missing data. In patients with cognitive impairment, informed consent was obtained from the health care proxy, and the CGA was administered with the health care proxy. Therefore, denominators are less than the 337 patients who were randomized in each group. Health care proxies were not always able to indicate whether patients had a certain geriatric condition.

^b Score range of 0 to 5, with increased risk of functional impairment at a score of 2 or higher.¹⁹

^c Range of 0 to 6, with a higher score indicating more disabilities.²¹

^d Range of 0 to 31, with a higher score indicating more or more severe comorbidity.²⁴

^e Vascular, neurodegenerative, musculoskeletal, or endocrine diagnosis at admission.

^f Use of 5 or more different medications.

^g One question to the patient: "Do you experience incontinence for urine or stool?"

^h One question to the patient: "Do you have an indwelling urinary catheter?"

ⁱ One question to the patient: "Do you experience dizziness?"

^j Score of 2 or higher on the Short Nutritional Assessment Questionnaire (range, 0-7).²²

^k Score of 4 or higher on the Visual Analogue Pain Intensity Scale (range, 0-10).²⁸

^l Cognitively impaired if a score of less than 24 on the Mini-Mental State Examination (range, 0-30).²⁰

^m Both questions on the 2-item Geriatric Depression Scale.²⁹

ⁿ One question to the patient: "Did you experience a fall during the past 6 months?"

^o One question to the patient: "Do you have trouble falling asleep or sleeping through the night?"


^p One question to the patient: "Do you have a hearing impairment, regardless of the use of a hearing device?"

^q One question to the patient: "Do you have a visual impairment, regardless of the use of eyeglasses?"

of the participants. Seventy-three percent (199 of 272) of participants had a CCRN transitional care bridge visit during hospitalization. The first home visit was conducted for

89.7% (244 of 272) of the participants within 48 hours after discharge. There was no difference in the total number of visits in the intervention arm between those discharged to a

Figure 2. Mortality From Admission to 6 Months After Admission



Shown are survival curves from a Cox proportional hazards regression analysis adjusted for study site and cognitive functioning (Mini-Mental State Examination Score, <24 vs ≥24), with the numbers of participants at risk for each 30-day interval for the intervention arm and the CGA-only arm. Significant protective intervention effects were observed for 1-month mortality and 6-month mortality. CGA indicates comprehensive geriatric assessment.

Table 3. Differences Between the Intervention Arm and the CGA-Only Control Arm for ADL and Cognition

Variable	LS Mean (95% CI)		CGA Only		P Value for Treatment × Time Interaction
	Intervention	6 Months	Baseline	6 Months	
ADL ^a	1.81 (1.63-1.99)	2.00 (1.78-2.23)	1.89 (1.71-2.07)	1.92 (1.69-2.15)	.32
ADL7 ^b	1.83 (1.59-2.07)	3.66 (3.40-3.92)	1.92 (1.68-2.16)	3.80 (3.54-4.06)	.78
MMSE ^c	22.2 (21.3-23.0)	24.8 (23.8-25.8)	22.4 (21.6-23.3)	25.1 (24.1-26.1)	.87

Abbreviations: ADL, activities of daily living; CGA, comprehensive geriatric assessment; LS, least squares; MMSE, Mini-Mental State Examination.

^a As measured with the 6-item Katz Index of ADL.²¹ Estimates are the adjusted LS means (95% CIs) for the number of ADL that were disabled at 2 weeks before hospitalization and at 6 months after admission (see the Statistical Analysis subsection of the Methods section) using a linear mixed-effects model with random participant-specific intercepts and center and stratification factor MMSE score (<24 vs ≥24).

^b Represents a sensitivity analysis in which decedents were assigned a score of 7 on the Katz Index of ADL.

^c Cognitive functioning, measured with the 11-item MMSE.²⁰ Estimates are the adjusted LS means (95% CIs) for the MMSE score at the time of hospital admission and at 6 months after admission (see the Statistical Analysis subsection of the Methods section) using a linear mixed-effects model with random participant-specific intercepts and center.

nursing home (mean [SD], 3.8 [1.7] visits) and those directly discharged home (mean [SD], 3.9 [1.7] visits).

Activities of Daily Living

There was no difference in the mean Katz Index of ADL at 6 months between the intervention arm (mean, 2.0; 95% CI, 1.8-2.2) and the CGA-only arm (mean, 1.9; 95% CI, 1.7-2.2). For sensitivity analysis, decedents were assigned a score of higher-than-maximum disability, yet no difference in the number of ADL disabilities was detected between the intervention arm (3.6; 95% CI, 3.4-3.92) and the control arm (3.8; 95% CI, 3.5-4.1) (P = .78). The ADLs at 2 weeks before hospitalization for participants who died (mean [SD], 2.2 [1.9]) and participants who survived (mean [SD], 1.6 [1.6]) to 6 months differed significantly (P = .002).

Mortality

There were 85 deaths (25.2%) in the intervention arm and 104 deaths (30.9%) in the CGA-only arm. Figure 2 shows the survival curve until 6 months after admission. Significant protective intervention effects were observed for 1-month mortality (hazard ratio [HR], 0.63; 95% CI, 0.39-0.99; P = .047) and 6-month mortality (HR, 0.75; 95% CI, 0.56-0.99; P = .045). The

number needed to treat to prevent one death was 16. The intervention started once the CCRN visited the participant in the hospital (median, 4 days after admission); therefore, we conducted a sensitivity analysis with 2 intervention participants and 3 CGA-only participants removed because they died before the first visit. The effects of the intervention at 1 month (HR, 0.63; 95% CI, 0.39-1.01; P = .05) and at 6 months (HR, 0.75; 95% CI, 0.56-1.00; P = .05) did not reach statistical significance.

Readmission, Institutionalization, and Cognitive Functioning

There were 106 readmissions in the intervention arm (33.5% of 316 discharged from the hospital) and 88 readmissions in the CGA-only arm (29.0% of 303 discharged from the hospital). No effect of the intervention was seen on the time to the first unplanned readmission by 6 months (HR, 1.21; 95% CI, 0.91-1.60; P = .76).

The numbers of discharges to a nursing home were 51 in the intervention arm (16.1% of 316 discharged alive) and 41 in the CGA-only arm (13.5% of 303 discharged alive). Among those discharged to a nursing home, the time to discharge home did not significantly differ, with a median of 63 days (IQR, 27-138 days) for the intervention arm vs a median of 38 days (IQR,

16-76 days) for the CGA-only arm ($P = .76$). Similarly, there was no effect of the intervention on cognitive functioning at 6 months after discharge ($P = .87$) (Table 3).

Subgroup Analysis

For 6-month mortality, there were no significant interactions between the assigned treatment and either the prespecified subgroups (ISAR-HP score of 2-3 vs 4-5) or the Charlson Comorbidity Index (0-3 vs ≥ 4). These results are shown in eTable 3 in Supplement 2.

Discussion

In this multicenter randomized clinical trial in acutely hospitalized older participants comparing systematic CGA followed by a transitional care program with systematic CGA only, we found no effect of the intervention on ADL disability at 6 months after discharge. For the secondary outcome analysis, we observed lower 1-month and 6-month mortality rates among participants in the intervention arm. The intervention had no effect on all other secondary outcomes of cognitive functioning, the time to unplanned hospital readmission, or the time to discharge from a nursing home to the community by 6 months after discharge.

While we did not observe any differences between the intervention and CGA-only arms in the change in ADLs, only a few transitional care trials have included ADL functioning as an outcome. Naylor and colleagues¹⁴ found a short-term improvement in functioning in an intervention arm who received transitional care, but this effect was not found at 6 months. We only measured ADL functioning at 6 months after discharge; therefore, we could not have observed early improvements in functioning. Another possible explanation of the lack of an intervention effect may be that, because both arms received CGA during hospitalization, both arms' health care professionals were aware of the risk of functional decline. Furthermore, we had broad inclusion criteria, which may have increased the variability of response to the intervention, and we provided a structured exercise intervention.

Our finding of a reduction of the intervention on mortality at 1 month and 6 months after discharge should be taken with caution because the primary outcome and all other secondary outcomes of this study were negative. However, other studies of similar interventions have suggested an effect on mortality. Deschodt et al⁸ showed that the systematic CGA approach by a geriatric consultation team led to a reduction in

6-month mortality. Other transitional care trials often start at the time of hospital admission but do not include a CGA or an engagement during hospitalization by a geriatric consultation team.^{14,15,30} Naylor and colleagues¹⁴ conducted a trial in patients with heart failure, including a geriatric assessment and follow-up by an advanced practice nurse, which found a reduction in the combined end point of readmission or mortality. Other transitional care trials either have not studied mortality as an end point³⁰⁻³⁴ or have not found reductions in mortality.^{15,16}

In contrast with other studies^{12,35} on transitional care, we did not observe a reduction in unplanned readmissions in the intervention arm. We observed lower 6-month readmission rates compared with other trials that have been conducted.¹² We hypothesize that these lower rates may result from the high standard of primary health care in the Dutch health care system, with easily accessible general practitioner care and home care for all citizens.

There are some limitations to our study. Despite the efforts to complete the follow-up assessments, we had missing ADL outcome data among survivors. However, our analytic model included all randomized persons in both the random intercept and baseline measure. The 6-month outcome assessment closely followed the 6-month visit by the CCRN. As a result, some of the participants considered the additional home visit for the outcome assessment to be burdensome. Moreover, the outcome assessments of the participants with cognitive impairment had to be conducted with the closest proxy, and some proxies did not reply to requests for assessment. The strength of the present trial is the inclusion of vulnerable patients with a high risk of functional decline as well as those with cognitive impairment. These groups are often excluded from trials.

Conclusions

This multicenter randomized clinical trial on systematic CGA and transitional care until 6 months after discharge demonstrated no effect of the intervention on ADL functioning compared with systematic CGA alone. Although there was a significant reduction in mortality at 1 month and 6 months after admission, there were no effects on other secondary outcomes. A systematic CGA, followed by a transitional care program, might improve patient safety during the vulnerable period that occurs shortly after hospital discharge. Further studies are needed to confirm these results.

ARTICLE INFORMATION

Accepted for Publication: December 6, 2015.

Published Online: February 15, 2016.
doi:10.1001/jamainternmed.2015.8042.

Author Affiliations: Section of Geriatric Medicine, Department of Internal Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands (Buurman, Parlevliet, de Rooij); Department of Internal Medicine, Yale School of Medicine, New Haven, Connecticut (Buurman, Allore); Department of Biostatistics, Yale

School of Public Health, New Haven, Connecticut (Allore); Department of Internal Medicine, Onze Lieve Vrouwe Gasthuis, Amsterdam, the Netherlands (Blok); Department of Internal Medicine, Flevo Hospital, Almere, the Netherlands (van Deelen); Department of Geriatric Medicine, Kennemer Gasthuis, Haarlem, the Netherlands (van Deelen); Department of General Practice, University of Amsterdam, Amsterdam, the Netherlands (Moll van Charante); Clinical Research Unit, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands

(de Haan); Department of Internal Medicine, University Medical Center Groningen, Groningen, the Netherlands (de Rooij).

Author Contributions: Drs Buurman and Parlevliet contributed equally to the article and share first authorship. Dr Buurman had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Buurman, Parlevliet,

Blok, van Deelen, Moll van Charante, de Haan, de Rooij.
Acquisition, analysis, or interpretation of data: All authors.
Drafting of the manuscript: Buurman, Parlevliet, Allore, de Rooij.
Critical revision of the manuscript for important intellectual content: All authors.
Statistical analysis: Buurman, Allore.
Obtained funding: de Rooij.
Administrative, technical, or material support: All authors.
Study supervision: Buurman, Parlevliet, Blok, van Deelen, de Haan, de Rooij.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study was funded by ZonMW, the Netherlands Organization for Health Research and Development, project number 311020201. Dr Buurman was supported by a Rubicon grant from the Netherlands Organization of Health Research (NWO), 825.12.022, Nederlands Trial Register NTR2384.

Role of the Funder/Sponsor: The funding sources had no role in the design and conduct of the study; the data collection, management, analysis, or interpretation of the data; the drafting or review of the manuscript; and the submission of the manuscript.

Additional Contributions: Mark Trentalange, MD, MPH, from the Program on Aging at the Yale School of Medicine, provided support with the data analyses. We thank all the members of the geriatric consultation teams at the participating hospitals, the community care registered nurses who provided the intervention in the home care setting, and all the older patients who participated in the trial.

REFERENCES

- Buurman BM, Hoogerduijn JG, de Haan RJ, et al. Geriatric conditions in acutely hospitalized older patients: prevalence and one-year survival and functional decline. *PLoS One*. 2011;6(11):e26951. doi:10.1371/journal.pone.0026951.
- Gill TM, Allore HG, Gahbauer EA, Murphy TE. Change in disability after hospitalization or restricted activity in older persons. *JAMA*. 2010;304(17):1919-1928.
- Boyd CM, Landefeld CS, Counsell SR, et al. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. *J Am Geriatr Soc*. 2008;56(12):2171-2179.
- Kansagara D, Englander H, Salanitro A, et al. Risk prediction models for hospital readmission: a systematic review. *JAMA*. 2011;306(15):1688-1698.
- Walter LC, Brand RJ, Counsell SR, et al. Development and validation of a prognostic index for 1-year mortality in older adults after hospitalization. *JAMA*. 2001;285(23):2987-2994.
- Ellis G, Whitehead MA, Robinson D, O'Neill D, Langhorne P. Comprehensive geriatric assessment for older adults admitted to hospital: meta-analysis of randomised controlled trials. *BMJ*. 2011;343:d6553.
- Naylor MD, Aiken LH, Kurtzman ET, Olds DM, Hirschman KB. The care span: the importance of transitional care in achieving health reform. *Health Aff (Millwood)*. 2011;30(4):746-754.
- Deschodt M, Flamaing J, Haentjens P, Boonen S, Milisen K. Impact of geriatric consultation teams on clinical outcome in acute hospitals: a systematic review and meta-analysis. *BMC Med*. 2013;11:48.
- Kripalani S, Roumie CL, Dalal AK, et al; PILL-CVD (Pharmacist Intervention for Low Literacy in Cardiovascular Disease) Study Group. Effect of a pharmacist intervention on clinically important medication errors after hospital discharge: a randomized trial. *Ann Intern Med*. 2012;157(1):1-10.
- Kripalani S, LeFevre F, Phillips CO, Williams MV, Basaviah P, Baker DW. Deficits in communication and information transfer between hospital-based and primary care physicians: implications for patient safety and continuity of care. *JAMA*. 2007;297(8):831-841.
- Coleman EA, Chugh A, Williams MV, et al. Understanding and execution of discharge instructions. *Am J Med Qual*. 2013;28(5):383-391.
- Verhaegh KJ, MacNeil-Vroomen JL, Eslami S, Geerlings SE, de Rooij SE, Buurman BM. Transitional care interventions prevent hospital readmissions for adults with chronic illnesses. *Health Aff (Millwood)*. 2014;33(9):1531-1539.
- Prvu Bettger J, Alexander KP, Dolor RJ, et al. Transitional care after hospitalization for acute stroke or myocardial infarction: a systematic review. *Ann Intern Med*. 2012;157(6):407-416.
- Naylor MD, Brooten DA, Campbell RL, Maislin G, McCauley KM, Schwartz JS. Transitional care of older adults hospitalized with heart failure: a randomized, controlled trial [published correction appears in *J Am Geriatr Soc*. 2004;52(7):1228]. *J Am Geriatr Soc*. 2004;52(5):675-684.
- Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med*. 2006;166(17):1822-1828.
- Parry C, Min SJ, Chugh A, Chalmers S, Coleman EA. Further application of The Care Transitions Intervention: results of a randomized controlled trial conducted in a fee-for-service setting. *Home Health Care Serv Q*. 2009;28(2-3):84-99.
- Boter H, van Delden JJ, de Haan RJ, Rinkel GJ. Modified informed consent procedure: consent to postponed information. *BMJ*. 2003;327(7409):284-285.
- Buurman BM, Parlevliet JL, van Deelen BA, de Haan RJ, de Rooij SE. A randomised clinical trial on a comprehensive geriatric assessment and intensive home follow-up after hospital discharge: the transitional care bridge. *BMC Health Serv Res*. 2010;10:296.
- Hoogerduijn JG, Buurman BM, Korevaar JC, Grobbee DE, de Rooij SE, Schuurmans MJ. The prediction of functional decline in older hospitalised patients. *Age Ageing*. 2012;41(3):381-387.
- Folstein MF, Folstein SE, McHugh PR. "Mini-Mental State": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975;12(3):189-198.
- Katz S, Ford AB, Moskowitz RW, Jackson BA, Jaffe MW. Studies of illness in the aged: the Index of ADL: a standardized measure of biological and psychosocial function. *JAMA*. 1963;185(12):914-919.
- Kruizenga HM, Seidel JC, de Vet HC, Wierdsma NJ, van Bokhorst-de van der Schueren MA. Development and validation of a hospital screening tool for malnutrition: the Short Nutritional Assessment Questionnaire (SNAQ). *Clin Nutr*. 2005;24(1):75-82.
- Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: the confusion assessment method: a new method for detection of delirium. *Ann Intern Med*. 1990;113(12):941-948.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383.
- Altman DG, Andersen PK. Calculating the number needed to treat for trials where the outcome is time to an event. *BMJ*. 1999;319(7223):1492-1495.
- Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc*. 1999;94(446):496-509.
- Hochberg Y. A sharper Bonferroni procedure of multiple tests of significance. *Biometrika*. 1988;75(4):800-802.
- Collins SL, Moore RA, McQuay HJ. The Visual Analogue Pain Intensity Scale: what is moderate pain in millimetres? *Pain*. 1997;72(1-2):95-97.
- Yesavage JA, Sheikh JI. Geriatric Depression Scale (GDS): recent evidence and development of a shorter version. *Clin Gerontologist*. 1986;5(1):165-173.
- Kwok T, Lee J, Woo J, Lee DT, Griffith S. A randomized controlled trial of a community nurse-supported hospital discharge programme in older patients with chronic heart failure. *J Clin Nurs*. 2008;17(1):109-117.
- Naylor MD, Brooten D, Campbell R, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA*. 1999;281(7):613-620.
- Jack BW, Chetty VK, Anthony D, et al. A reengineered hospital discharge program to decrease rehospitalization: a randomized trial. *Ann Intern Med*. 2009;150(3):178-187.
- Saleh SS, Freire C, Morris-Dickinson G, Shannon T. An effectiveness and cost-benefit analysis of a hospital-based discharge transition program for elderly Medicare recipients. *J Am Geriatr Soc*. 2012;60(6):1051-1056.
- Zhao Y, Wong FK. Effects of a postdischarge transitional care programme for patients with coronary heart disease in China: a randomised controlled trial. *J Clin Nurs*. 2009;18(17):2444-2455.
- Leppin AL, Gionfriddo MR, Kessler M, et al. Preventing 30-day hospital readmissions: a systematic review and meta-analysis of randomized trials. *JAMA Intern Med*. 2014;174(7):1095-1107.